

Section 2.3: Infinite sets and cardinality

Recall from Section 2.2 that

- The cardinality of a finite set is defined as the number of elements in it.
- Two sets A and B have **the same cardinality** if (and only if) it is possible to match each element of A to an element of B in such a way that every element of each set has exactly one “partner” in the other set.

In the case of finite sets, the second point above might seem to be overcomplicating the issue, since we can tell if two finite sets have the same cardinality by just counting their elements and noting that they have the same number.

Two sets have **the same cardinality** if (and only if) it is possible to match each element of A to an element of B in such a way that every element of each set has exactly one “partner” in the other set.

The notion of bijective correspondence is emphasized for two reasons.

- It is occasionally possible to establish that two finite sets are in bijective correspondence without knowing the cardinality of either of them.
- We can't count the number of elements in an infinite set. However, for a given pair of infinite sets, we could possibly show that it is or isn't possible to construct a bijective correspondence between them.

Infinite sets having the same cardinality

Definition

Suppose that A and B are sets (finite or infinite). We say that A and B have the same cardinality (written $|A| = |B|$) if a bijective correspondence exists between A and B .

In other words, A and B have the **same cardinality** if it's possible to match each element of A to a different element of B in such a way that every element of both sets is matched exactly once. In order to say that A and B have **different** cardinalities we need to establish that it's **impossible** to match up their elements with a bijective correspondence. If A and B are infinite sets, showing that such a thing is *impossible* can be a formidable challenge.

$$\mathbb{Z} = \{ \dots, -2, -1, 0, \underbrace{1, 2, 3, \dots} \}$$


Definition

The set \mathbb{N} of natural numbers (“counting numbers”) consists of all the positive integers. $\mathbb{N} = \{1, 2, 3, \dots\}$.

Example

Show that \mathbb{N} and \mathbb{Z} have the same cardinality.


We need to fill the right-hand column of the table below with the integers *in some order*, in such a way that each integer appears there exactly once.

\mathbb{N}		\mathbb{Z}	
1	\longleftrightarrow	?	
2	\longleftrightarrow	?	
3	\longleftrightarrow	?	
4	\longleftrightarrow	?	
\vdots	\longleftrightarrow	\vdots	

Bijjective correspondence between \mathbb{N} and \mathbb{Z}

So we need to list all the integers on the right hand side, in such a way that every integer appears once. Just following the natural order on the integers won't work, because then there is no first entry for our list.

\mathbb{N}		\mathbb{Z}
1	\longleftrightarrow	?
2	\longleftrightarrow	?
3	\longleftrightarrow	?
4	\longleftrightarrow	?
\vdots	\longleftrightarrow	\vdots



Starting at a particular integer like 0 and then following the natural order won't work, because then we will never get (for example) any negative integers in our list.


\mathbb{N}		\mathbb{Z}
1	\longleftrightarrow	?
2	\longleftrightarrow	?
3	\longleftrightarrow	?

Bijective correspondence $\mathbb{N} \longleftrightarrow \mathbb{Z}$

We can start with 0, then list 1 and then -1 , then 2 and then -2 , then 3 and then -3 and so on. This is a systematic way of writing out **all** the integers, in which each appears exactly once. Our table becomes

\mathbb{N}		\mathbb{Z}
1	\longleftrightarrow	0 ✓
2	\longleftrightarrow	1 ✓
3	\longleftrightarrow	-1 ✓
4	\longleftrightarrow	2
5	\longleftrightarrow	-2
6	\longleftrightarrow	3
\vdots	\longleftrightarrow	\vdots

22



Exercise 41


*What integer corresponds to the natural number 22 in the list?
In what position does the integer -63 appear?*


from Lecture 12:

A pair of infinite sets have the same cardinality if there is a bijjective correspondence between them - i.e. its possible to match every element of each set with an element of the other one, so that every element of each set has exactly one "companion" in the other.

Example \mathbb{N} and \mathbb{Z}
 $\{1, 2, 3, 4, \dots\}$ $\{\dots, -2, -1, 0, 1, 2, 3, 4, \dots\}$

↓ ↓ ↓ ↓
? ? ? ?
-3





A more explicit version



If we want to be fully explicit about how this bijective correspondence works, we can even give a formula for the integer that is matched to each natural number. The correspondence above describes a bijective function $f : \mathbb{N} \longrightarrow \mathbb{Z}$ given for $n \in \mathbb{N}$ by

$$f(n) = \begin{cases} \frac{n}{2} & \text{if } n \text{ is even} \\ -\left(\frac{n-1}{2}\right) & \text{if } n \text{ is odd} \end{cases}$$

As well as understanding this example at the informal/intuitive level suggested by the picture above, think about the formula above, and satisfy yourself that it does indeed describe a bijection between \mathbb{N} and \mathbb{Z} .


Curiosities of Infinite Sets

The example above demonstrates a curious thing that can happen when considering cardinalities of infinite sets. The set \mathbb{N} of natural numbers is a proper subset of the the set \mathbb{Z} of integers (this means that every natural number is an integer, but the natural numbers do not account for all the integers).

Yet we have just shown that \mathbb{N} and \mathbb{Z} can be put in bijective correspondence. So it is possible for an infinite set to be in bijective correspondence with a proper subset of itself, and hence to have the same cardinality as a proper subset of itself.

This can't happen for finite sets (why?).

A and B finite sets $A \subsetneq B$
 $|A| < |B|$ e.g. A might have 5 elements and B might have 20



Countably Infinite Sets

Putting an infinite set in bijective correspondence with \mathbb{N} amounts to providing a robust and unambiguous scheme or instruction for listing all its elements starting with a first, then a second, third, etc., in such a way that it can be seen that every element of the set will appear exactly once in the list.

Definition

A set is called countably infinite (or denumerable) if it can be put in bijective correspondence with the set of natural numbers. A set is called countable if it is either finite or countably infinite.


Basically, an infinite set is countable if its elements can be listed in an inclusive and organised way. “Listable” might be a better word, but it is not really used.

Countability of \mathbb{Q} ← fractions a/b (a, b integers)

Thus the sets \mathbb{N} and \mathbb{Z} have the same cardinality. Maybe this is not so surprising, because these sets have a strong geometric resemblance as sets of points on the number line.

What is more surprising is that \mathbb{N} (and hence \mathbb{Z}) has the same cardinality as the set \mathbb{Q} of all **rational** numbers. These sets do not resemble each other much geometrically. The natural numbers are **sparse** and **evenly spaced**, whereas the rational numbers are **densely packed** into the number line.

Nevertheless, as the following construction shows, \mathbb{Q} is a countable set.



\mathbb{Q} is countable

We need to show that the rational numbers can be organized into a numbered list in a systematic way that includes all of them. Such a list is a one-to-correspondence with the set \mathbb{N} of natural numbers.


Start with the following array of fractions.

...	$-3/1$	$-2/1$	$-1/1$	$0/1$	$1/1$	$2/1$	$3/1$...
...	$-3/2$	$-2/2$	$-1/2$	$0/2$	$1/2$	$2/2$	$3/2$...
...	$-3/3$	$-2/3$	$-1/3$	$0/3$	$1/3$	$2/3$	$3/3$...
...	$-3/4$	$-2/4$	$-1/4$	$0/4$	$1/4$	$2/4$	$3/4$...
	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	

\mathbb{Q} is countable

We need to show that the rational numbers can be organized into a numbered list in a systematic way that includes all of them. Such a list is a one-to-correspondence with the set \mathbb{N} of natural numbers.

Construct a path through the whole array :



\mathbb{Q} is countable

In these fractions, the numerators increase through all the integers as we travel along the rows, and the denominators increase through all the natural numbers as we travel downwards through the columns.

Every rational number occurs somewhere in the array.

This path determines a listing of all the fractions in the array, that starts as follows

0/1, 1/1, 1/2, 0/2, -1/2, -1/1, -2/1, -2/2, -2/3, -1/3, 0/3, 1/3, 2/3

\mathbb{Q} is countable

$0/1, 1/1, 1/2, 0/2, -1/2, -1/1, -2/1, -2/2, -2/3, -1/3, 0/3, 1/3,$
 $2/3, 2/2, 2/1, 3/1, 3/2, 3/3, 3/4, \dots$

What this construction demonstrates is a bijective correspondence between the set \mathbb{N} of natural numbers and the set of all fractions in our array.

This is not (exactly) a bijective correspondence between \mathbb{N} and \mathbb{Q} .

Exercise 42

Why not? (Think about this before reading on.)

The reason why not is that every rational number appears many times in our array.

\mathbb{Q} is countable

In order to get a bijective correspondence between \mathbb{N} and \mathbb{Q} , construct a list of all the rational numbers from the array as above, but whenever a rational number is encountered that has already appeared, leave it out. Our list will begin

$$0/1, 1/1, 1/2, -1/2, -1/1, -2/1, -2/3, -1/3, 1/3, 2/3, 2/1, \\ 3/1, 3/2, 3/4, \dots$$

We conclude that the rational numbers are countable.

Note : Unlike our one-to-one correspondence between \mathbb{N} and \mathbb{Z} , in this case we cannot write down a simple formula to tell us what rational number will be Item 34 on our list (i.e. corresponds to the natural number 34) or where in our list the rational number $292/53$ will appear.