

We could forget about the third element of S and just write

$$\begin{pmatrix} a \\ b \end{pmatrix} = (a + b) \begin{pmatrix} 3 \\ 1 \end{pmatrix} + (-a) \begin{pmatrix} 2 \\ 1 \end{pmatrix}.$$

So the three elements of S are not necessary to form a spanning set of \mathbb{R}^2 . We could span \mathbb{R}^2 just with the subset $\{\begin{pmatrix} 2 \\ 1 \end{pmatrix}, \begin{pmatrix} 3 \\ 1 \end{pmatrix}\}$ of S . We note that $\begin{pmatrix} 1 \\ -1 \end{pmatrix}$ is a \mathbb{R} -linear combination of the other two elements of S . If we drop this element from S , we can still recover it in the span of the remaining elements.

The second example above motivates this lemma, which we will explore further in the next section.

Lemma 2.1.4. *Suppose that $S_1 \subset S$, where S is a subset of a vector space V . Then $\langle S_1 \rangle \subseteq \langle S \rangle$, and $\langle S_1 \rangle = \langle S \rangle$ if and only if every element of $S \setminus S_1$ is a linear combination of elements of S_1 .*

We finish this section by noting the distinction between a *finite dimensional* and *infinite dimensional* vector space.

Definition 2.1.5. *A vector space is said to be finite dimensional if it has a finite spanning set. A vector space that does not have a finite spanning set is infinite dimensional.*

Two examples of infinite dimensional vector spaces

1. The vector space $\mathbb{R}[x]$ of all polynomials with real coefficients is infinite dimensional. To see this, let S be any finite subset of $\mathbb{R}[x]$ (i.e. a finite set of polynomials). Let x^k be the highest power of x to appear in any element of S . Then no linear combination of elements of S has degree exceeding x^k , so the linear span of S cannot be all of $\mathbb{R}[x]$.
2. The set \mathbb{R} of real numbers is infinite dimensional as a vector space over the field \mathbb{Q} of rational numbers.

2.2 Linear Independence

Definition 2.2.1. *Let S be a subset of a vector space V , having at least 2 elements. Then S is linearly independent if no element of S is a linear combination of the other elements of S (equivalently, if no element of S belongs to the span of the other elements of S).*

A subset consisting of a single element is linear independent, provided that its unique element is not the zero vector.

To decide if a given set is linearly independent, Definition 2.2.1 is maybe not the most useful formulation, because it requires us to check something separately for each element of S , which could take a lot of work. The following alternative version of the definition might have less appeal at an intuitive level, but it is often more useful in practice.

Definition 2.2.2. *Let S be a non-empty subset of a vector space V . Then S is linearly independent if the only way to write the zero vector in V as a linear combination of elements of S is to take all the coefficients to be 0.*

Equivalence of the two definitions

Let $S = \{v_1, \dots, v_k\}$ and suppose that $v_1 \in \langle v_2, \dots, v_k \rangle$. Then

$$v_1 = a_2 v_2 + \dots + a_k v_k,$$

and

$$0 = -v_1 + a_2 v_2 + \dots + a_k v_k$$

is an expression for the zero vector as a linear combination of elements of S , whose coefficients are not all zero.

On the other hand, suppose that

$$0 = c_1 v_1 + \cdots + c_k v_k,$$

where the scalars c_i are not all zero. If $c_1 \neq 0$ (for example), then the above equation can be rearranged to express v_1 as a linear combination of v_2, \dots, v_k :

$$v_1 = -\frac{c_2}{c_1} v_2 - \cdots - \frac{c_k}{c_1} v_k.$$

Example 2.2.3. In \mathbb{R}^3 , let $S = \left\{ \begin{bmatrix} 1 \\ 2 \\ -1 \end{bmatrix}, \begin{bmatrix} -2 \\ 3 \\ 2 \end{bmatrix}, \begin{bmatrix} -3 \\ 8 \\ 3 \end{bmatrix} \right\}$.

To determine whether S is linearly independent, we must investigate whether the system of equations

$$x \begin{bmatrix} 1 \\ 2 \\ -1 \end{bmatrix} + y \begin{bmatrix} -2 \\ 3 \\ 2 \end{bmatrix} + z \begin{bmatrix} -3 \\ 8 \\ 3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

has solutions other than $(x, y, z) = (0, 0, 0)$. The augmented matrix of this system, and its RREF, are

$$\begin{bmatrix} 1 & -2 & -3 & 0 \\ 2 & 3 & 8 & 0 \\ -1 & 2 & 3 & 0 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & 1 & 0 \\ 0 & 1 & 2 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Thus for any t , $(x, y, z) = (-t, -2t, t)$ is a solution, and for example by taking $t = 1$ we see that

$$-1 \begin{bmatrix} 1 \\ 2 \\ -1 \end{bmatrix} - 2 \begin{bmatrix} -2 \\ 3 \\ 2 \end{bmatrix} + 1 \begin{bmatrix} -3 \\ 8 \\ 3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix},$$

and hence that each of the three elements of S is a linear combination of the other two. So S is not linearly independent (we say that S is *linearly dependent*).

We note below some characterizations of linearly independent set. Let S be a subset of a vector space V .

1. S is linearly independent if S is a *minimal* spanning set of its linear span - no proper subset of S spans the same subspace of V that S does, or every proper subspace of S spans a strictly smaller subspace than S itself.
2. S is linearly independent if every element of $\langle S \rangle$ has a *unique* expression as a linear combination of elements of S . If a particular element of $\langle S \rangle$ had two different expressions as linear combinations of S , with different coefficients, then subtracting one from the other would give a non-trivial expression for 0_V as a linear combination of elements of S , and by Definition 2.2.2 we would conclude that S is linearly dependent.
3. Another version of 2. above: S is linearly independent if every element of the span of S has *unique coordinates* in terms of the elements of S .

So a linearly independent set in a vector space V is a *minimal* or *irredundant* spanning set for its linear span. If its linear span happens to be all of V , it gets a special name.

Definition 2.2.4. A basis of a vector space V is a spanning set of V that is linearly independent.