

# Lecture 16: Eigenvectors and Diagonalizability

March 8, 2024

# Lecture 16: Eigenvectors and Diagonalizability

- 1 Representing a linear transformation with respect to different bases
- 2 Diagonalizability
- 3 Non-diagonalizability

## Example from Lecture 15

Let  $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$  be the linear transformation defined by  $v \rightarrow Av$ , where  $A = \begin{bmatrix} -2 & 2 & 1 \\ 4 & 5 & -1 \\ -4 & -8 & 3 \end{bmatrix}$ . The matrix of  $T$  with respect to the (ordered) basis  $B$  of  $\mathbb{R}^3$  with elements  $b_1 = \begin{bmatrix} 1 \\ 0 \\ 4 \end{bmatrix}$ ,  $b_2 = \begin{bmatrix} 2 \\ -1 \\ 0 \end{bmatrix}$ ,  $b_3 = \begin{bmatrix} 0 \\ -1 \\ 2 \end{bmatrix}$  is

$$\begin{bmatrix} 2 & 0 & 0 \\ 0 & -3 & 0 \\ 0 & 0 & 7 \end{bmatrix}.$$

This means:  $T(b_1) = 2b_1$ ,  $T(b_2) = 3b_2$ ,  $T(b_3) = 7b_3$ , and for any  $v \in \mathbb{R}^3$ ,

$$\underbrace{[T(v)]_B}_{B\text{-coordinates of } T(v)} = \underbrace{A'[v]_B}_{\text{matrix-vector product}}.$$

## Change of basis again

Let  $P$  be the matrix with the basis vectors from  $B$  as columns.

From Lecture 14,  $P^{-1}$  is the **change of basis matrix** from the standard basis to  $B$ . For any element  $v$  of  $\mathbb{R}^3$ , its  $B$ -coordinates are given by the matrix-vector product

$$[v]_B = P^{-1}v.$$

Equivalently, if we start with the  $B$ -coordinates, then the standard coordinates of  $v$  are given by

$$v = P[v]_B.$$

So  $P$  itself is the change of basis matrix from  $B$  to the standard basis.

## Similarity - the relation of $A$ and $A'$

Starting with  $A$ , the matrix of  $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$  with respect to the **standard basis**, how do we find  $A'$  the matrix of  $T$  with respect to  $B$ ?

- 1 Take a vector  $v$  of  $\mathbb{R}^3$ , written in  $B$ -coordinates as the column  $[v]_B$ .
- 2 Convert to standard coordinates (so that we can apply  $T$  by multiplying by  $A$ ): take the product  $P[v]_B$ .
- 3 Apply  $T$ : left-multiply by  $A$  to get  $AP[v]_B$ . This column has the **standard coordinates** of  $T(v)$ .
- 4 Convert to  $B$ -coordinates: left-multiply by  $P^{-1}$ , the change of basis matrix from standard to  $B$ . This gives  $P^{-1}AP[v]_B$ . This column has the  **$B$ -coordinates** of  $T(v)$ .
- 5 Conclusion: For any element  $v$  of  $\mathbb{R}^3$ , the  $B$ -coordinates of  $T(v)$  are given by  $(P^{-1}AP)[v]_B$ .

The  **$B$ -matrix of  $T$**  is  $P^{-1}AP$ , where  $P$  has the elements of  $B$  as columns.

# Similar Matrices

**Definition** Two square matrices  $A$  and  $B$  are **similar** if  $B = P^{-1}AP$  for an invertible matrix  $P$ .

## Notes

- 1 Two distinct matrices are similar if and only if they represent the same linear transformation, with respect to different bases.
- 2 We can't tell by glancing at a pair of square matrices if they are similar or not, but there is one feature that is easy to check. The **trace** of a square matrix is the sum of the entries on the main diagonal, from top left to bottom right. If two matrices are similar, they have the same trace.
- 3 Similar matrices also have some other features in common, such as having the same determinant.
- 4 Our example showed that the  $3 \times 3$  matrix  $A = \begin{bmatrix} -2 & 2 & 1 \\ 4 & 5 & -1 \\ -4 & -8 & 3 \end{bmatrix}$  is similar to the **diagonal** matrix  $\text{diag}(2, -3, 7)$ . We say  $A$  is **diagonalizable** in this situation.

# Two equivalent interpretations of diagonalizability

- 1 From the diagonal form of  $A'$  we have  $T(b_1) = 2b_1$ ,  $T(b_2) = -3b_2$  and  $T(b_3) = 7b_3$ . This means that each of the basis elements  $b_1, b_2, b_3$  is mapped by  $T$  to a scalar multiple of itself - each of them is an *eigenvector* of  $T$ .
- 2 We can rearrange the version  $P^{-1}AP = A'$  to  $AP = PA'$ . Bearing in mind that  $P = \begin{bmatrix} | & | & | \\ b_1 & b_2 & b_3 \\ | & | & | \end{bmatrix}$  and that  $A' = \text{diag}(2, -3, 7)$ , this is saying that

$$A \begin{bmatrix} | & | & | \\ b_1 & b_2 & b_3 \\ | & | & | \end{bmatrix} = \begin{bmatrix} | & | & | \\ b_1 & b_2 & b_3 \\ | & | & | \end{bmatrix} \begin{bmatrix} 2 & 0 & 0 \\ 0 & -3 & 0 \\ 0 & 0 & 7 \end{bmatrix} \Rightarrow \begin{bmatrix} | & | & | \\ Ab_1 & Ab_2 & Ab_3 \\ | & | & | \end{bmatrix} = \begin{bmatrix} | & | & | \\ 2b_1 & -3b_2 & 7b_3 \\ | & | & | \end{bmatrix}$$

This means that  $Ab_1 = 2b_1$ ,  $Ab_2 = -3b_2$  and  $Ab_3 = 7b_3$ , so that  $B = \{b_1, b_2, b_3\}$  is a basis of  $\mathbb{R}^3$  consisting of *eigenvectors* of  $A$ .

# Eigenvectors and Diagonalizability

**Definition** An **eigenvector** of a square matrix  $A$  is a non-zero column vector  $v$  for which  $Av = \lambda v$  for some scalar  $\lambda$ , called the **eigenvalue** of  $A$  to which  $v$  corresponds.

The **eigenvalues** of  $A$  are the roots of its **characteristic polynomial**  $\det(\lambda I_n - A)$ .

The **eigenspace** corresponding to a particular eigenvalue  $\lambda$  is the set of all vectors  $v$  satisfying  $Av = \lambda v$ . It is a subspace of the relevant  $\mathbb{R}^n$ , of dimension at least 1.

The matrix  $A \in M_n(\mathbb{R})$  is **diagonalizable** if and only if  $\mathbb{R}^n$  has a basis consisting of eigenvectors of  $A$ . In this case  $P^{-1}AP$  is diagonal, where  $P$  is a matrix whose  $n$  columns are linearly independent eigenvectors of  $A$ . The diagonal entries of  $P^{-1}AP$  are the corresponding eigenvalues.



# Non-diagonalizability (two examples)

For  $A \in M_n(\mathbb{R})$ , it does not always happen that  $\mathbb{R}^n$  has a basis consisting of eigenvectors of  $A$ .

## Examples

- 1 The matrix  $A = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$  is diagonalizable in  $M_2(\mathbb{C})$  but not in  $M_2(\mathbb{R})$ .

This matrix represents a clockwise **rotation through  $90^\circ$**  about the origin. It does not fix any line in  $\mathbb{R}^2$ . Its characteristic polynomial is  $\lambda^2 + 1$ .

- 2 The matrix  $B = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$  is not diagonalizable even over  $\mathbb{C}$ .

This matrix represents a **horizontal shear**. Its characteristic polynomial is  $(\lambda - 1)^2$  but its 1-eigenspace consists only of the  $X$ -axis. It does not have two linearly independent eigenvectors.