

Lecture 10: Subspaces and Spanning Sets

February 17, 2025

Lecture 10: Linear Transformations and Subspaces

- 1 Matrix multiplication is composition of functions
- 2 The kernel and image
- 3 Subspaces
- 4 Spanning sets
- 5 Spanning Sets

Matrix multiplication is composition

Suppose that $T : \mathbb{R}^n \rightarrow \mathbb{R}^p$ and $S : \mathbb{R}^p \rightarrow \mathbb{R}^m$ are linear transformations. Then $S \circ T$ (S after T) is the **linear transformation** from \mathbb{R}^n to \mathbb{R}^m defined for $v \in \mathbb{R}^n$ by

$$S \circ T(v) = S(T(v)).$$

Question How does the $(m \times n)$ matrix $M_{S \circ T}$ of $S \circ T$ depend on the $(m \times p)$ matrix M_S of S and the $(p \times n)$ matrix M_T of T ?

To answer this we have to think about the definition of $M_{S \circ T}$.

- Its first column has the coordinates of $S \circ T(e_1) = S(T(e_1))$.
- $T(e_1)$ is the first column of M_T .
- Then $S(T(e_1))$ is the matrix-vector product M_S [first column of M_T]. This is the first column of the matrix product $M_S M_T$.
- Same for all the other columns: the conclusion is $M_{S \circ T} = M_S M_T$.

Matrix multiplication is composition of linear transformations.

Corollary Matrix multiplication is associative.

The Image and Kernel of a Linear Transformation

$T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ is the linear transformation with $M_T = \begin{bmatrix} 1 & 2 & 0 \\ 2 & -1 & 5 \\ 1 & 1 & 1 \end{bmatrix}$.

The **image** of T is the subset of \mathbb{R}^3 consisting of all elements $T(v)$, where $v \in \mathbb{R}^3$. This is the set of all vectors of the form

$$a \begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix} + b \begin{bmatrix} 2 \\ -1 \\ 1 \end{bmatrix} + c \begin{bmatrix} 0 \\ 5 \\ 1 \end{bmatrix}.$$

In matrix terms, this is the **column space** of M_T .

The **kernel** of T is the set of all vectors v in \mathbb{R}^3 with $T(v) = 0$.

This is the set of all column vectors whose entries a, b, c satisfy

$$\begin{bmatrix} 1 & 2 & 0 \\ 2 & -1 & 5 \\ 1 & 1 & 1 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \end{bmatrix} = a \begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix} + b \begin{bmatrix} 2 \\ -1 \\ 1 \end{bmatrix} + c \begin{bmatrix} 0 \\ 5 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}.$$

In matrix terms this is the **(right) nullspace** of M_T .

Example: The kernel is a **line** and the image is a **plane**

$$\begin{bmatrix} 1 & 2 & 0 \\ 2 & -1 & 5 \\ 1 & 1 & 1 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} \Rightarrow \begin{bmatrix} 1 & 2 & 0 & | & 0 \\ 2 & -1 & 5 & | & 0 \\ 1 & 1 & 1 & | & 0 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & 2 & | & 0 \\ 0 & 1 & -1 & | & 0 \\ 0 & 0 & 0 & | & 0 \end{bmatrix}$$

The **kernel/nullspace** is $\boxed{\{(-2, 1, 1)t, t \in \mathbb{R}\}}$ a **line** in \mathbb{R}^3 .

That $(-2, 1, 1)$ is in the kernel of T means that (for example) Column 3 of M_T is a linear combination of Columns 1 and 2.

$$-2 \begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix} + 1 \begin{bmatrix} 2 \\ -1 \\ 1 \end{bmatrix} + 1 \begin{bmatrix} 0 \\ 5 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} \Rightarrow \begin{bmatrix} 0 \\ 5 \\ 1 \end{bmatrix} = 2 \begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix} - \begin{bmatrix} 2 \\ -1 \\ 1 \end{bmatrix}$$

It follows that every linear combination of all three columns of M_T is actually a linear combination just of Columns 1 and 2.

The column space of M_T is $\left\{ a \begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix} + b \begin{bmatrix} 2 \\ -1 \\ 1 \end{bmatrix} : a, b \in \mathbb{R} \right\}$, a **plane** in \mathbb{R}^3 .

Definition A (non-empty) subset V of \mathbb{R}^n is a **subspace** if

- It is **closed under addition**: $u + v \in V$ whenever $u \in V$ and $v \in V$.
- It is **closed under scalar multiplication**: $ku \in V$ whenever $u \in V$ and $k \in \mathbb{R}$.

Examples

- 1 $\{(x, y, z) \in \mathbb{R}^3 : x + y + z = 1\}$ is **not** a subspace of \mathbb{R}^3 . The vectors $(1, 0, 0)$ and $(0, 1, 0)$ belong to this set but their sum $(1, 1, 0)$ does not.
- 2 $\{(x, y, z) \in \mathbb{R}^3 : (x, y, z) \cdot (1, 2, 3) = 0\}$ **is** a subspace of \mathbb{R}^3 .
- 3 $\{(x, y, z) \in \mathbb{R}^3 : (x, y, z) \cdot (1, 2, 3) \neq 0\}$ is **not** a subspace of \mathbb{R}^3 . For example, $(1, 4, 1)$ and $(-5, -2, -1)$ belong to this set but their sum $(-4, 2, 0)$ does not.
- 4 The kernel of any linear transformation is a subspace.
- 5 The image of any linear transformation is a subspace.

Exercise Prove these last two points.

How to make subspaces

Let $S = \{v_1, \dots, v_k\}$ be any (finite) subset of \mathbb{R}^n .

The subset of \mathbb{R}^n consisting of all linear combinations of the elements of S is a subspace of \mathbb{R}^n , denoted by $\langle S \rangle$ or $\langle v_1, v_2, \dots, v_k \rangle$ and called the linear span (or just span) of S .

Proof (that $\langle S \rangle$ is a subspace).

Closed under **addition**: let $u, v \in \langle S \rangle$. Then $u = a_1 v_1 + a_2 v_2 + \dots + a_k v_k$, and $v = c_1 v_1 + c_2 v_2 + \dots + c_k v_k$, where the a_i and b_i are scalars. We need to show that $u + v \in \langle S \rangle$, which means showing that it is a linear combination of v_1, \dots, v_k . This is straightforward after everything has been set up, since $u + v = (a_1 + c_1)v_1 + (a_2 + c_2)v_2 + \dots + (a_k + c_k)v_k$. So S is closed under addition.

Closed under **scalar multiplication**: let $u \in \langle S \rangle$ and $c \in \mathbb{R}$. We need to show that cu is a linear combination of v_1, \dots, v_k . We know that $u = a_1 v_1 + a_2 v_2 + \dots + a_k v_k$, for scalars a_1, \dots, a_k . Then $cu = ca_1 v_1 + ca_2 v_2 + \dots + ca_k v_k$, so $cu \in \langle S \rangle$.

Spanning Sets

Let V be a subspace of \mathbb{R}^n (possibly V is all of \mathbb{R}^n). A subset S of V is called a **spanning set** of V if $\langle S \rangle = V$.

This means that every element of V is a linear combination of the elements of S .

Example The set $\{e_1, e_2, e_3\}$ is a spanning set of \mathbb{R}^3 , where (as usual)

$$e_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \quad e_2 = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \quad e_3 = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}. \quad \text{This is saying that every}$$

element of \mathbb{R}^3 is a linear combination of e_1, e_2, e_3 . For example

$$\begin{bmatrix} 2 \\ -3 \\ 4 \end{bmatrix} = 2e_1 - 3e_2 + 4e_3.$$

Remark A set S of three column vectors in \mathbb{R}^3 is a **spanning set** of \mathbb{R}^3 if and only if each of e_1, e_2, e_3 is a linear combination of elements of S .

This occurs **if and only if** the 3×3 matrix whose columns are the vectors in S has an *inverse*.

Questions about Spanning Sets

- 1 Does \mathbb{R}^3 have a spanning set with fewer than three elements?
- 2 Does every spanning set of \mathbb{R}^3 have exactly three elements?
NO (why not?)
- 3 Does every spanning set of \mathbb{R}^3 **contain** one with exactly three elements?
- 4 If V is a **subspace** of \mathbb{R}^3 , does V have a spanning set with at most three elements?
- 5 If V is a **proper subspace** of \mathbb{R}^3 (i.e. not all of \mathbb{R}^3) does V have a spanning set with fewer than three elements?

Note A pair of vectors in \mathbb{R}^3 (if they are not scalar multiples of each other) span a **plane**. Adding a third vector (if it does not lie in this plane) gives a spanning set for all of \mathbb{R}^3 .